

$$I_{G,k} = \langle g_1, \dots, g_n \rangle$$

# Combinatorial Commutative Algebra, Graph Colorability, and Algorithms


Chris Hillar  
(University of California, Berkeley)

Joint with Troels Windfeldt  
(University of Copenhagen)

# Outline of talk

- Introduction: Colorability
- Motivational Problem
- Previous Work:  $k$ -colorability
  - algebraic characterizations
  - Groebner bases
- Notation and Color Encodings
- Statement of Main Results
- Algorithms

# Graph Colorings



Let  $G$  be a simple graph  $G = (V, E)$   
with vertices  $V = \{1, 2, \dots, n\}$ , edges  $E$


**Def:** A  $k$ -coloring of  $G$  is an assignment  
of  $k$  colors to the vertices of  $G$

**Def:** A  $k$ -coloring is **proper** if adjacent  
vertices receive different colors

**Def:** A graph is  $k$ -colorable if it has a proper  
 $k$ -coloring

# Unique Colorability

A 3-colorable graph that is not 2-colorable:




Other than permuting the colors, no other proper 3-colorings

**Def:** A **uniquely  $k$ -colorable** graph is one which has exactly  $k!$  proper  $k$ -colorings. In this case, the coloring is unique up to permutation of colors

# Technical Observation

$G$  cannot be uniquely  $k$ -colorable if there is a proper  $k$ -coloring **not** using all  $k$  colors



Not uniquely 3-colorable **although** there are  $3! = 6$  proper 3-colorings of  $G$  that use all 3 colors

# Motivational Problem


In 1990, Xu showed that uniquely  $k$ -colorable graphs must have many edges

**Theorem** (Xu): If  $G$  is uniquely  $k$ -colorable then

$$|E| \geq (k-1)n - k(k-1)/2$$

In example,  $7 \geq 2 \cdot 5 - 3 = 7$

Notice:  $G$  contains a **triangle**



# Xu's Conjecture

In general, he conjectured that if equality holds,  $G$  must contain a  $k$ -clique.

**Conjecture (Xu):** If  $G$  is uniquely  $k$ -colorable and  $|E| = (k-1)n - k(k-1)/2$ , then  $G$  contains a  $k$ -clique

This conjecture was shown to be false in 2001 by Akbari et al using a combinatorial argument

We wanted to find a **JPE proof** (Just Press Enter)


# Computer Proof

This leads to the following concrete problem:

**Problem:** Find an (effective) algorithm to decide unique  $k$ -colorability (and find the coloring).

3-colorable?

Uniquely 3-colorable?






# Computer Proof

This leads to the following concrete problem:

**Problem:** Find an (effective) algorithm to decide unique  $k$ -colorability (and find the coloring).

3-colorable?

Uniquely 3-colorable?



# Computer Proof


This leads to the following concrete problem:

**Problem:** Find an (effective) algorithm to decide unique  $k$ -colorability (and find the coloring).

3-colorable?

Uniquely 3-colorable?

Y e s !



# Computer Proof

This leads to the following concrete problem:


**Problem:** Find an (effective) algorithm to decide unique  $k$ -colorability (and find the coloring).

3-colorable?

Uniquely 3-colorable?

Y e s !

Y e s !



# Computer Proof

This leads to the following concrete problem:

**Problem:** Find an (effective) algorithm to decide unique  $k$ -colorability (and find the coloring).


3-colorable?

Uniquely 3-colorable?

Y e s !

Y e s !


. . .



# Counterexample to Xu

How about the graph actually used


by Akbari, Mirrokni, Sadjad:  $|E| = 45$ ,  $n = 24$ ,  $k = 3$




# Algebraic Colorability

**Main Idea** (implicit in work of Bayer, de Loera, Lovász):

colorings are points in varieties



varieties are represented by ideals



ideals can be manipulated with **Groebner Bases**

# $k$ -Colorings as Points in Varieties

**Setup:**  $F$  is an algebraically closed field,  $(\text{char } F) \nmid k$   
So  $F$  contains  $k$  distinct  $k$ th roots of unity. Let

$$I_k = \langle x_1^k - 1, x_2^k - 1, \dots, x_n^k - 1 \rangle \subset F[x_1, \dots, x_n]$$


This ideal is radical, and  $|V(I_k)| = k^n$

---

Can think of point  $v = (v_1, \dots, v_n)$  in  $V(I_k)$  as **assignment**

$$v = (v_1, \dots, v_n) \iff \text{vertex } i \text{ gets color } v_i$$

Eg. If  $1 = \text{Blue}$ ,  $-1 = \text{Red}$ , then  $v = (1, -1, 1)$  is coloring



# Proper $k$ -Colorings of Graphs

We can also restrict to proper  $k$ -colorings of graph  $G$

$$I_{G,k} = I_k + \langle x_i^{k-1} + x_i^{k-2}x_j + \cdots + x_j^{k-1} : (i,j) \in E \rangle$$

This ideal is radical, and  $|V(I_{G,k})| = \#$  proper  $k$ -colorings

---

Proof: ( $\Leftarrow$ ) If  $v$  proper, then  $v_i \neq v_j$  and

$$(v_i - v_j) \cdot (v_i^{k-1} + \cdots + v_j^{k-1}) = v_i^k - v_j^k = 1 - 1 = 0$$

Thus,  $v_i^{k-1} + \cdots + v_j^{k-1} = 0$  and  $v$  in  $V(I_{G,k})$




# Algebraic Characterization

Notice that if  $I_{G,k} = \langle 1 \rangle = F[x_1, \dots, x_n]$  then

$$V(I_{G,k}) = \emptyset \Rightarrow G \text{ is not } k\text{-colorable}$$

Therefore, we have a **test for  $k$ -colorability**:

**Algorithm:** Compute a reduced **Groebner basis**  $B$  for  $I_{G,k}$ .  
Then,  $B = \{1\}$  iff  $G$  is **not**  $k$ -colorable.



$(k = 2)$

$$I_{G,k} = \langle x_1^2 - 1, x_2^2 - 1, x_3^2 - 1, \\ x_1 + x_2, x_2 + x_3, x_1 + x_3 \rangle \\ = \langle 1 \rangle$$

$$2x_1^2 = (x_1 - x_2)(x_1 + x_2) + (x_2 - x_3)(x_2 + x_3) + (x_1 - x_3)(x_1 + x_3)$$

# Graph Polynomial


The **graph polynomial** of  $G$  encodes adjacency of vertices **algebraically**:

$$f_G = \prod_{\substack{\{i,j\} \in E \\ i < j}} (x_i - x_j)$$

One should think of  $f_G$  as a **test polynomial** for  $k$ -colorability.

$$I_3 = \langle x_1^3 - 1, x_2^3 - 1, x_3^3 - 1 \rangle$$

$$k = 3, n = 3$$



Notice that  $f_G = (x_1 - x_2)(x_2 - x_3)(x_1 - x_3) \notin I_k$

**Conclusion:** There is a 3-coloring that is proper!

# Characterization Theorem

**$k$ -colorability Theorem** (Bayer, de Loera, Alon, Tarsi, Mnuk, Kleitman, Lovász): The following are equivalent

- (1)  $G$  is not  $k$ -colorable
- (2)  $I_{G,k} = \langle 1 \rangle$
- (3)  $f_G$  is contained in  $I_k$  (colorings zero  $f_G$ )

**Corollary:** There are simple tests for  $k$ -colorability involving polynomial algebra.

**Our goal:** Develop a similar **characterization theorem** for **unique  $k$ -colorability** and give a complete description of  $I_{G,k}$  when  $G$  is  $k$ -colorable

# Preparation: Color classes

Given  $G$  which is  $k$ -colorable with a coloring  $v = (v_1, \dots, v_n)$  using all  $k$  colors, we define:

Color class of  $i = \text{cl}(i) = \{ j : v_j = v_i \}$

representative of  $\text{cl}(i) = \max\{ j : j \text{ is in } \text{cl}(i) \}$

Denote these representatives


$$m_1 < m_2 < \dots < m_k = n$$

---

Eg.  $\text{cl}(1) = \{1, 5\}$ ,  $\text{cl}(2) = \{2, 3\}$

$\text{cl}(4) = \{4\}$

$$m_1 = 3 < m_2 = 4 < m_3 = 5$$



# New Polynomial Encoding

We need a **replacement** for the graph polynomial  $f_G$  in the statement of the  $k$ -colorability theorem

**Def:** Let  $U \subseteq \{1, \dots, n\}$ . Then we set  $h_U^d$  to be the **sum of all monomials of degree  $d$**  in the variables  $\{x_i : i \text{ in } U\}$

Eg.  $U = \{1, 2, 3\}$ ,  $d = 2$

$$h_U^d = x_1^2 + x_2^2 + x_3^2 + x_1x_2 + x_1x_3 + x_2x_3$$

Also, we set  $h_U^0 = 1$

# Replacements for $f_G$

Def: Given a proper  $k$ -coloring  $v$ , for each vertex  $i$ , let

$$g_i = \begin{cases} x_i^k - 1 & i = m_k (= n) \\ h^j_{\{m_j, \dots, m_k\}} & i = m_j \text{ for some } j \neq k \\ h^1_{\{i, m_2, \dots, m_k\}} & i \text{ in } \text{cl}(m_1) \\ x_i - x_{\max \text{ cl}(i)} & \text{otherwise} \end{cases}$$


$$g_5 = x_5^3 - 1$$

$$g_4 = h^2_{\{4,5\}} = x_4^2 + x_4 x_5 + x_5^2$$

$$g_3 = h^1_{\{3,4,5\}} = x_3 + x_4 + x_5$$

$$g_2 = h^1_{\{2,4,5\}} = x_2 + x_4 + x_5$$

$$g_1 = x_1 - x_5 \quad (m_1, m_2, m_3) = (3, 4, 5)$$



# Can go backwards

Find a proper  $k$ -coloring giving a set

$$g_i = \begin{cases} x_i^k - 1 & i = m_k (= n) \\ h^j_{\{m_j, \dots, m_k\}} & i = m_j \text{ for some } j \neq k \\ h^1_{\{i, m_2, \dots, m_k\}} & i \text{ in } \text{cl}(m_1) \\ x_i - x_{\max \text{ cl}(i)} & \text{otherwise} \end{cases}$$


$$g_5 = x_5^3 - 1$$

$$g_4 = h^2_{\{4,5\}} = x_4^2 + x_4 x_5 + x_5^2$$

$$g_3 = h^1_{\{3,4,5\}} = x_3 + x_4 + x_5$$

$$g_2 = h^1_{\{2,4,5\}} = x_2 + x_4 + x_5$$

$$g_1 = x_1 - x_5 \quad (m_1, m_2, m_3) = (?, ?, ?)$$



# Can go backwards

Find a proper  $k$ -coloring giving a set

$$g_i = \begin{cases} x_i^k - 1 & i = m_k (= n) \\ h^j_{\{m_j, \dots, m_k\}} & i = m_j \text{ for some } j \neq k \\ h^1_{\{i, m_2, \dots, m_k\}} & i \text{ in } \text{cl}(m_1) \\ x_i - x_{\max \text{cl}(i)} & \text{otherwise} \end{cases}$$


$$g_5 = x_5^3 - 1$$

$$g_4 = h^2_{\{4,5\}} = x_4^2 + x_4 x_5 + x_5^2$$

$$g_3 = h^1_{\{3,4,5\}} = x_3 + x_4 + x_5$$

$$g_2 = h^1_{\{2,4,5\}} = x_2 + x_4 + x_5$$

$$g_1 = x_1 - x_5 \quad (m_1, m_2, m_3) = (?, ?, 5)$$





# Can go backwards

Find a proper  $k$ -coloring giving a set

$$g_i = \begin{cases} x_i^k - 1 & i = m_k (= n) \\ h^j_{\{m_j, \dots, m_k\}} & i = m_j \text{ for some } j \neq k \\ h^1_{\{i, m_2, \dots, m_k\}} & i \text{ in } \text{cl}(m_1) \\ x_i - x_{\max \text{cl}(i)} & \text{otherwise} \end{cases}$$


$$g_5 = x_5^3 - 1$$

$$g_4 = h^2_{\{4,5\}} = x_4^2 + x_4 x_5 + x_5^2$$

$$g_3 = h^1_{\{3,4,5\}} = x_3 + x_4 + x_5$$

$$g_2 = h^1_{\{2,4,5\}} = x_2 + x_4 + x_5$$

$$g_1 = x_1 - x_5 \quad (m_1, m_2, m_3) = (?, 4, 5)$$



# Can go backwards

Find a proper  $k$ -coloring giving a set

$$g_i = \begin{cases} x_i^k - 1 & i = m_k (= n) \\ h^j_{\{m_j, \dots, m_k\}} & i = m_j \text{ for some } j \neq k \\ h^1_{\{i, m_2, \dots, m_k\}} & i \text{ in } \text{cl}(m_1) \\ x_i - x_{\max \text{cl}(i)} & \text{otherwise} \end{cases}$$


$$g_5 = x_5^3 - 1$$

$$g_4 = h^2_{\{4,5\}} = x_4^2 + x_4 x_5 + x_5^2$$

$$g_3 = h^1_{\{3,4,5\}} = x_3 + x_4 + x_5$$

$$g_2 = h^1_{\{2,4,5\}} = x_2 + x_4 + x_5$$

$$g_1 = x_1 - x_5 \quad (m_1, m_2, m_3) = (3, 4, 5)$$



# Can go backwards

Find a proper  $k$ -coloring giving a set

$$g_i = \begin{cases} x_i^k - 1 & i = m_k (= n) \\ h^j_{\{m_j, \dots, m_k\}} & i = m_j \text{ for some } j \neq k \\ h^1_{\{i, m_2, \dots, m_k\}} & i \text{ in } \text{cl}(m_1) \\ x_i - x_{\max \text{ cl}(i)} & \text{otherwise} \end{cases}$$


$$g_5 = x_5^3 - 1$$

$$g_4 = h^2_{\{4,5\}} = x_4^2 + x_4 x_5 + x_5^2$$

$$g_3 = h^1_{\{3,4,5\}} = x_3 + x_4 + x_5$$

$$g_2 = h^1_{\{2,4,5\}} = x_2 + x_4 + x_5$$

$$g_1 = x_1 - x_5 \quad (m_1, m_2, m_3) = (3, 4, 5)$$



# Can go backwards

Find a proper  $k$ -coloring giving a set

$$g_i = \begin{cases} x_i^k - 1 & i = m_k (= n) \\ h^j_{\{m_j, \dots, m_k\}} & i = m_j \text{ for some } j \neq k \\ h^1_{\{i, m_2, \dots, m_k\}} & i \text{ in } \text{cl}(m_1) \\ x_i - x_{\max \text{ cl}(i)} & \text{otherwise} \end{cases}$$


$$g_5 = x_5^3 - 1$$

$$g_4 = h^2_{\{4,5\}} = x_4^2 + x_4 x_5 + x_5^2$$

$$g_3 = h^1_{\{3,4,5\}} = x_3 + x_4 + x_5$$

$$g_2 = h^1_{\{2,4,5\}} = x_2 + x_4 + x_5$$

$$g_1 = x_1 - x_5 \quad (m_1, m_2, m_3) = (3, 4, 5)$$



Encoding:  $v \dashrightarrow A_v = \langle g_1, \dots, g_n \rangle$

The set of polynomials  $\{g_1, \dots, g_n\}$  encodes the coloring  $v$

**Lemma:** Let  $A_v = \langle g_1, \dots, g_n \rangle$

- (1)  $I_{G,k} \subseteq A_v$
- (2)  $A_v$  is a radical ideal
- (3)  $|V(A_v)| = k!$

**Interpretation:**

- (1) zeroes of  $A_v$  are proper  $k$ -colorings of  $G$
- (2)  $A_v$  and its zeroes are in 1-1 correspondence
- (3) Up to permutation,  $A_v$  encodes precisely  $v$

# Characterization Theorem

**Theorem** [-,W 06]: Let  $G$  be a graph. The following are equivalent

(1)  $G$  is  $k$ -colorable

(2)  $\bigcap A_v \subseteq I_{G,k}$

(3)  $\bigcap A_v = I_{G,k}$

**Point:** We have found an interpretation involving ideals for the statement:

$$\bigcup \{ V(A_v) : v \text{ is proper} \} = \{ \text{proper colorings} \}$$

# Unique Characterization

In general, the map from proper  $k$ -colorings

$$v \mapsto A_v = \langle g_1, \dots, g_n \rangle$$

only depends on how  $v$  partitions  $V = \{1, \dots, n\}$  into color classes. In particular,

**Fact:** If  $G$  is uniquely colorable, then there is a unique set of polynomials  $\{g_1, \dots, g_n\}$  that corresponds to all  $v$ .

**Proof:** All  $v$  partition  $V$  the same way

# Unique Characterization

**Corollary** [-,W 06]: Fix a **proper  $k$ -coloring**  $v$  of  $G$ . Let  $A_v = \langle g_1, \dots, g_n \rangle$ . The following are equivalent

- (1)  $G$  is uniquely  $k$ -colorable
- (2)  $g_1, \dots, g_n$  belong to  $I_{G,k}$
- (3)  $A_v = \langle g_1, \dots, g_n \rangle = I_{G,k}$

More canonically, we have the following

**Theorem** [-,W 06]:  $G$  is **uniquely  $k$ -colorable** if and only if the **reduced Groebner basis** for  $I_{G,k}$  (w.r.t any term order with  $x_n < \dots < x_1$ ) has the form  $g_1, \dots, g_n$



# Algorithms

The main theorems give algorithms for determining unique colorability


**Algorithm 1:** Given a proper  $k$ -coloring, construct the polynomials  $g_1, \dots, g_n$  and reduce them modulo  $I_{G,k}$

**Algorithm 2:** Compute the reduced Groebner basis for  $I_{G,k}$  and see whether it has the form  $g_1, \dots, g_n$ ; if so, read off the coloring.

- (easy to check form and to read off coloring)

# Just Press Enter

Computing with a field  $F$  with  $\text{char } F = 2$ , we find that (after pressing enter and waiting 5 seconds) the following graph is indeed *uniquely  $k$ -colorable*.



# The End

(of talk)