Harmonic Functions on $\mathbb{R}^n$

Justin Corvino

Lafayette College
U.S.A.

June 10, 2012

Introduction

We will use conformal changes of metric to deform the scalar curvature, and so we will want to understand the behavior of the Laplace operator Δ_g, as for $u > 0$:

$$R(u^{4/(n-2)}g) = -\frac{4(n-1)}{n-2}u^{-\frac{n+2}{n-2}}(\Delta_g u - \frac{n-2}{4(n-1)}R(g)u).$$

Introduction

We will use conformal changes of metric to deform the scalar curvature, and so we will want to understand the behavior of the Laplace operator Δ_g, as for $u > 0$:

$$R(u^{4/(n-2)}g) = -\frac{4(n-1)}{n-2}u^{-\frac{n+2}{n-2}}(\Delta_g u - \frac{n-2}{4(n-1)}R(g)u).$$

Recall that the Hessian of a function on a Riemannian (or Lorentzian) manifold (M, g) is $\nabla(du)$, the components of which are denoted

$$u_{;ij} = \nabla(du)(\partial_i, \partial_j) = (\nabla_{\partial_j} du)(\partial_i) = u_{,ij} - du(\nabla_{\partial_j}\partial_i) = u_{,ij} - \Gamma^k_{ij}u_{,k}.$$

Introduction

We will use conformal changes of metric to deform the scalar curvature, and so we will want to understand the behavior of the Laplace operator Δ_g, as for $u > 0$:

$$R(u^{4/(n-2)}g) = -\frac{4(n-1)}{n-2}u^{-\frac{n+2}{n-2}}(\Delta_g u - \frac{n-2}{4(n-1)}R(g)u).$$

Recall that the Hessian of a function on a Riemannian (or Lorentzian) manifold (M, g) is $\nabla(du)$, the components of which are denoted

$$u_{;ij} = \nabla(du)(\partial_i, \partial_j) = (\nabla_{\partial_j} du)(\partial_i) = u_{,ij} - du(\nabla_{\partial_j}\partial_i) = u_{,ij} - \Gamma^k_{ij}u_{,k}.$$

On a Riemannian manifold, we write $\Delta_g u = g^{ij}u_{;ij}$, which can also be written $\Delta_g u = \mathrm{div}_g(\mathrm{grad}_g u)$.
We often write $\Delta = \Delta_{g_E}$

On a Lorentzian manifold, the trace of the Hessian gives the *wave operator* $\Box_g$.

Introduction

- Laplace equation: $\Delta_g u = 0$ (u is *harmonic*)

Introduction

- Laplace equation: $\Delta_g u = 0$ (u is *harmonic*)
- Poisson equation: $\Delta_g u = f$

Introduction

- Laplace equation: $\Delta_g u = 0$ (u is *harmonic*)
- Poisson equation: $\Delta_g u = f$
- Eigenvalue problem: $\Delta_g u = -\lambda u$.

Introduction

- Laplace equation: $\Delta_g u = 0$ (u is *harmonic*)
- Poisson equation: $\Delta_g u = f$
- Eigenvalue problem: $\Delta_g u = -\lambda u$.

As $R(u^{4/(n-2)}g) = -\frac{4(n-1)}{n-2} u^{-\frac{n+2}{n-2}} (\Delta_g u - \frac{n-2}{4(n-1)} R(g) u)$,
if $g = g_E$ is Euclidean, then prescribing $R(u^{\frac{4}{n-2}} g) = 0$ is equivalent to $\Delta u = 0$, $R(u^{\frac{4}{n-2}} g) \geq 0$ is equivalent to $\Delta u \leq 0$ (u is *superharmonic*).

Introduction

- Laplace equation: $\Delta_g u = 0$ (u is *harmonic*)
- Poisson equation: $\Delta_g u = f$
- Eigenvalue problem: $\Delta_g u = -\lambda u$.

As $R(u^{4/(n-2)}g) = -\frac{4(n-1)}{n-2} u^{-\frac{n+2}{n-2}} (\Delta_g u - \frac{n-2}{4(n-1)} R(g) u)$,
if $g = g_E$ is Euclidean, then prescribing $R(u^{\frac{4}{n-2}} g) = 0$ is equivalent to $\Delta u = 0$, $R(u^{\frac{4}{n-2}} g) \geq 0$ is equivalent to $\Delta u \leq 0$ (u is *superharmonic*).

More generally, prescribing the scalar curvature of the conformal metric gives a semi-linear elliptic equation; prescribing that it vanish gives the linear equation $\Delta_g u - \frac{n-2}{4(n-1)} R(g) u = 0$.
So it is important to understand the behavior of operators of the form $(\Delta_g - f)$.

Introduction

Classical Newtonian gravity: $\mathbf{F} = -\frac{GmM}{r^2}\widehat{r}$, where $\mathbf{F}$ is the force on m, $\widehat{r}$ is the unit vector from M to m, and r is the distance from m to M.

Introduction

Classical Newtonian gravity: $\mathbf{F} = -\frac{GmM}{r^2}\widehat{r}$, where $\mathbf{F}$ is the force on m, $\widehat{r}$ is the unit vector from M to m, and r is the distance from m to M.

$\Phi = -\frac{GM}{|x|}$, where $|x|$ is the distance from M, is the *gravitational potential*.

$\Delta\Phi = 4\pi GM\delta_0$.

More generally, if σ is the mass density, $\Delta\Phi = 4\pi G\sigma$ determines the gravitational potential (along with boundary condition at infinity).

Introduction

Classical Newtonian gravity: $\mathbf{F} = -\frac{GmM}{r^2}\widehat{r}$, where $\mathbf{F}$ is the force on m, $\widehat{r}$ is the unit vector from M to m, and r is the distance from m to M.

$\Phi = -\frac{GM}{|x|}$, where $|x|$ is the distance from M, is the *gravitational potential*.

$\Delta\Phi = 4\pi GM\delta_0$.

More generally, if σ is the mass density, $\Delta\Phi = 4\pi G\sigma$ determines the gravitational potential (along with boundary condition at infinity).

Isolated gravitational system: If σ is compactly supported (or decays suitably at infinity), then how should we expect Φ to behave?

Spherically symmetric harmonic functions

On a Riemannian manifold (M, g), it is not a hard exercise to show

$$\Delta_g u = \frac{1}{\sqrt{\det g}} \frac{\partial}{\partial x^i} \left(\sqrt{\det g}\, g^{ij} \frac{\partial u}{\partial x^j} \right).$$

Spherically symmetric harmonic functions

On a Riemannian manifold (M, g), it is not a hard exercise to show

$$\Delta_g u = \frac{1}{\sqrt{\det g}} \frac{\partial}{\partial x^i} \left(\sqrt{\det g}\, g^{ij} \frac{\partial u}{\partial x^j} \right).$$

Now consider $\mathbb{R}^n$ with the Euclidean metric. If $u(x) = u(|x|)$ (slight abuse of notation), then with $r = |x|$, $g_E = dr^2 + r^{n-1} g_{\mathbb{S}^{n-1}}$, so

Spherically symmetric harmonic functions

On a Riemannian manifold (M, g), it is not a hard exercise to show

$$\Delta_g u = \frac{1}{\sqrt{\det g}} \frac{\partial}{\partial x^i} \left(\sqrt{\det g}\, g^{ij} \frac{\partial u}{\partial x^j} \right).$$

Now consider $\mathbb{R}^n$ with the Euclidean metric. If $u(x) = u(|x|)$ (slight abuse of notation), then with $r = |x|$, $g_E = dr^2 + r^{n-1} g_{\mathbb{S}^{n-1}}$, so

$$\Delta u = \frac{1}{r^{n-1}} \frac{d}{dr} \left(r^{n-1} \frac{du}{dr} \right) = u''(r) + \frac{n-1}{r} u'(r).$$

Spherically symmetric harmonic functions

On a Riemannian manifold (M, g), it is not a hard exercise to show

$$\Delta_g u = \frac{1}{\sqrt{\det g}} \frac{\partial}{\partial x^i} \left(\sqrt{\det g}\, g^{ij} \frac{\partial u}{\partial x^j} \right).$$

Now consider $\mathbb{R}^n$ with the Euclidean metric. If $u(x) = u(|x|)$ (slight abuse of notation), then with $r = |x|$, $g_E = dr^2 + r^{n-1} g_{\mathbb{S}^{n-1}}$, so

$$\Delta u = \frac{1}{r^{n-1}} \frac{d}{dr} \left(r^{n-1} \frac{du}{dr} \right) = u''(r) + \frac{n-1}{r} u'(r).$$

So $\Delta u = 0$ is equivalent to $0 = \frac{d}{dr} \left(r^{n-1} \frac{du}{dr} \right)$.

Spherically symmetric harmonic functions

On a Riemannian manifold (M, g), it is not a hard exercise to show

$$\Delta_g u = \frac{1}{\sqrt{\det g}} \frac{\partial}{\partial x^i} \left(\sqrt{\det g}\, g^{ij} \frac{\partial u}{\partial x^j} \right).$$

Now consider $\mathbb{R}^n$ with the Euclidean metric. If $u(x) = u(|x|)$ (slight abuse of notation), then with $r = |x|$, $g_E = dr^2 + r^{n-1} g_{\mathbb{S}^{n-1}}$, so

$$\Delta u = \frac{1}{r^{n-1}} \frac{d}{dr} \left(r^{n-1} \frac{du}{dr} \right) = u''(r) + \frac{n-1}{r} u'(r).$$

So $\Delta u = 0$ is equivalent to $0 = \frac{d}{dr} \left(r^{n-1} \frac{du}{dr} \right)$.
Integrating we obtain (A and B are constants)

$$u(r) = \begin{cases} Ar + B\,, & n = 1 \\ A \log r + B\,, & n = 2 \\ Ar^{2-n} + B\,, & n > 2 \end{cases}$$

Fundamental solution

We will stick to $n \geq 2$.

Fundamental solution

We will stick to $n \geq 2$.

Note that in dimension $n = 2$, $\log r$ is unbounded either above or below on $\mathbb{R}^2 \setminus \{0\}$, whereas $r^{2-n} > 0$ and goes to 0 as $|x| \to \infty$ for $n > 2$.

Fundamental solution

We will stick to $n \geq 2$.

Note that in dimension $n = 2$, $\log r$ is unbounded either above or below on $\mathbb{R}^2 \setminus \{0\}$, whereas $r^{2-n} > 0$ and goes to 0 as $|x| \to \infty$ for $n > 2$.

$n = 2$ special: harmonic functions tied in with holomorphic function theory.

Fundamental solution

We will stick to $n \geq 2$.

Note that in dimension $n = 2$, $\log r$ is unbounded either above or below on $\mathbb{R}^2 \setminus \{0\}$, whereas $r^{2-n} > 0$ and goes to 0 as $|x| \to \infty$ for $n > 2$.

$n = 2$ special: harmonic functions tied in with holomorphic function theory.

Fundamental Solution

As distributions,

$$\Delta\left(\frac{1}{2\pi}\log|x|\right) = \delta_0$$
$$\Delta\left(\frac{1}{(2-n)n\Omega_n}|x|^{2-n}\right) = \delta_0$$

$\Omega_n = \mathrm{vol}(\mathbb{B}^n(1))$ is the volume of the Euclidean unit ball. Note $n\Omega_n = \omega_{n-1}$ is the area of the unit sphere.

Fundamental solution

We will stick to $n \geq 2$.

Note that in dimension $n = 2$, $\log r$ is unbounded either above or below on $\mathbb{R}^2 \setminus \{0\}$, whereas $r^{2-n} > 0$ and goes to 0 as $|x| \to \infty$ for $n > 2$.

$n = 2$ special: harmonic functions tied in with holomorphic function theory.

Fundamental Solution

As distributions,

$$\Delta\left(\frac{1}{2\pi}\log|x|\right) = \delta_0$$
$$\Delta\left(\frac{1}{(2-n)n\Omega_n}|x|^{2-n}\right) = \delta_0$$

$\Omega_n = \mathrm{vol}(\mathbb{B}^n(1))$ is the volume of the Euclidean unit ball. Note $n\Omega_n = \omega_{n-1}$ is the area of the unit sphere.

Let $\Gamma(x,y) = \frac{1}{(2-n)n\Omega_n}|x-y|^{2-n}$ for $n > 2$, and $\Gamma(x,y) = \frac{1}{2\pi}\log|x-y|$ for $n = 2$.

Fundamental Solution

Fundamental Solution

What this means is that for $u \in C_c^2(\mathbb{R}^n)$,

$$u(x) = \begin{cases} \int_{\mathbb{R}^2} \frac{1}{2\pi} \log|x-y| \Delta u(y)\, dy\ , & n = 2 \\ \int_{\mathbb{R}^n} \frac{1}{(2-n)n\Omega_n} |x-y|^{2-n} \Delta u(y)\, dy\ , & n > 2 \end{cases}$$

Fundamental Solution

Fundamental Solution

What this means is that for $u \in C_c^2(\mathbb{R}^n)$,

$$u(x) = \begin{cases} \int_{\mathbb{R}^2} \frac{1}{2\pi} \log|x-y| \Delta u(y)\, dy\ , & n = 2 \\ \int_{\mathbb{R}^n} \frac{1}{(2-n)n\Omega_n} |x-y|^{2-n} \Delta u(y)\, dy\ , & n > 2 \end{cases}$$

Exercise

If $\Delta u = f$ compactly supported, $n > 2$, expand $|x-y|^{2-n}$ for large $|x|$ to obtain an expansion of u near infinity:

$$u(x) = \frac{A}{|x|^{n-2}} + \frac{x^i B_i}{|x|^n} + O(|x|^{-n}).$$

Find the constants A, B_i.

Mean Value Property

Again, we are discussing the Laplace operator on domains in Euclidean space.

Mean Value Property

Again, we are discussing the Laplace operator on domains in Euclidean space.

Mean Value Property

If $u \in C(\overline{B_R(a)})$ is harmonic on $\overline{B_R(a)}$, then

$$u(a) = \frac{1}{n\Omega_n R^{n-1}} \int\limits_{|x-a|=R} u(x)\, d\sigma = \frac{1}{\Omega_n R^n} \int\limits_{B_R(a)} u(x)\, dx.$$

Mean Value Property

Again, we are discussing the Laplace operator on domains in Euclidean space.

Mean Value Property

If $u \in C(\overline{B_R(a)})$ is harmonic on $\overline{B_R(a)}$, then

$$u(a) = \frac{1}{n\Omega_n R^{n-1}} \int\limits_{|x-a|=R} u(x)\, d\sigma = \frac{1}{\Omega_n R^n} \int\limits_{B_R(a)} u(x)\, dx.$$

Proof: Apply Green's identity $\int_\Omega (u\Delta v - v\Delta u)\, dx = \int_{\partial\Omega} (u\frac{\partial v}{\partial n} - v\frac{\partial u}{\partial n})\, d\sigma$ with $v = |x-a|^{2-n}$ (in case $n > 2$, and analogously in case $n = 2$), with $\Omega = \{x : \epsilon < |x-a| < R\}$. Let $\epsilon \to 0^+$ and use continuity.

Maximum Principle

MVP implies Maximum Principle

If u is harmonic on Ω a connected domain, then if u attains a max or min inside Ω, then u is constant.

Maximum Principle

MVP implies Maximum Principle

If u is harmonic on Ω a connected domain, then if u attains a max or min inside Ω, then u is constant.

If Ω bounded, u is harmonic on Ω and continuous on $\overline{\Omega}$, then the max and min of u are attained on $\partial\Omega$.

Dirichlet problem on a ball

Let $B_R = B_R(0)$. Let $P(x,y) = \frac{R^2 - |x|^2}{n\Omega_n R|x-y|^n}$. Then for $x \in B_R$, $\int_{y \in \partial B_R} P(x,y)\, d\sigma_y = 1$.

Dirichlet problem on a ball

Let $B_R = B_R(0)$. Let $P(x,y) = \frac{R^2 - |x|^2}{n\Omega_n R|x-y|^n}$. Then for $x \in B_R$, $\int_{y \in \partial B_R} P(x,y)\, d\sigma_y = 1$.

Dirichlet problem

If $\varphi \in C(\partial B_R)$, then $u(x) = \int_{y \in \partial B_R} \varphi(y) P(x,y)\, d\sigma_y$ is harmonic on B_R, continuous on $\overline{B_R}$, and $u = \varphi$ on ∂B_R.

Poisson equation on a ball

For any $y \in B_R$, solve $\Delta h_y(x) = 0$ on B_R with $h_y(x) = -\Gamma(x, y)$ for $x \in \partial B_R$. Let $G(x, y) = \Gamma(x, y) + h_y(x)$.

Poisson equation on a ball

For any $y \in B_R$, solve $\Delta h_y(x) = 0$ on B_R with $h_y(x) = -\Gamma(x, y)$ for $x \in \partial B_R$. Let $G(x, y) = \Gamma(x, y) + h_y(x)$.

Fact: $P = \frac{\partial G}{\partial n} = \frac{\partial G}{\partial |x|}$ on ∂B_R.

Poisson equation on a ball

For any $y \in B_R$, solve $\Delta h_y(x) = 0$ on B_R with $h_y(x) = -\Gamma(x, y)$ for $x \in \partial B_R$. Let $G(x, y) = \Gamma(x, y) + h_y(x)$.

Fact: $P = \frac{\partial G}{\partial n} = \frac{\partial G}{\partial |x|}$ on ∂B_R.

Applying Green's identity, and the distributional Laplacian of Γ (and hence of the *Green's function* G), we have

Poisson equation on ball

$$u(x) = \int_{\partial B_R} \varphi(y) P(x, y)\, d\sigma_y + \int_{B_R} G(x, y) f(y)\, dy$$

solves $\Delta u = f$ on B_R, $u = \varphi$ on ∂B_R.

Removable Singularity Theorem

Theorem

An isolated singularity of a bounded harmonic function is removable.

Removable Singularity Theorem

Theorem

An isolated singularity of a bounded harmonic function is removable.

Proof: WLOG, u is bounded and harmonic on $\overline{B} \setminus \{0\}$, where $B = B_1(0)$.

Removable Singularity Theorem

Theorem

An isolated singularity of a bounded harmonic function is removable.

Proof: WLOG, u is bounded and harmonic on $\overline{B} \setminus \{0\}$, where $B = B_1(0)$. Let v be harmonic on B, continuous on $\overline{B}$, with the same boundary values as u.

Removable Singularity Theorem

Theorem

An isolated singularity of a bounded harmonic function is removable.

Proof: WLOG, u is bounded and harmonic on $\overline{B} \setminus \{0\}$, where $B = B_1(0)$. Let v be harmonic on B, continuous on $\overline{B}$, with the same boundary values as u.
For $n > 2$, let $v_\epsilon(x) = u(x) - v(x) + \epsilon(|x|^{2-n} - 1)$.

Removable Singularity Theorem

Theorem

An isolated singularity of a bounded harmonic function is removable.

Proof: WLOG, u is bounded and harmonic on $\overline{B} \setminus \{0\}$, where $B = B_1(0)$.
Let v be harmonic on B, continuous on $\overline{B}$, with the same boundary values as u.
For $n > 2$, let $v_\epsilon(x) = u(x) - v(x) + \epsilon(|x|^{2-n} - 1)$.
Then $v_\epsilon = 0$ on ∂B, while (u and v are bounded) $v_\epsilon(x) \to +\infty$ as $x \to 0$.

Removable Singularity Theorem

Theorem

An isolated singularity of a bounded harmonic function is removable.

Proof: WLOG, u is bounded and harmonic on $\overline{B} \setminus \{0\}$, where $B = B_1(0)$.
Let v be harmonic on B, continuous on $\overline{B}$, with the same boundary values as u.
For $n > 2$, let $v_\epsilon(x) = u(x) - v(x) + \epsilon(|x|^{2-n} - 1)$.
Then $v_\epsilon = 0$ on ∂B, while (u and v are bounded) $v_\epsilon(x) \to +\infty$ as $x \to 0$.
By MP, $v_\epsilon(x) \geq 0$ on $B \setminus \{0\}$.

Removable Singularity Theorem

Theorem

An isolated singularity of a bounded harmonic function is removable.

Proof: WLOG, u is bounded and harmonic on $\overline{B} \setminus \{0\}$, where $B = B_1(0)$.
Let v be harmonic on B, continuous on $\overline{B}$, with the same boundary values as u.
For $n > 2$, let $v_\epsilon(x) = u(x) - v(x) + \epsilon(|x|^{2-n} - 1)$.
Then $v_\epsilon = 0$ on ∂B, while (u and v are bounded) $v_\epsilon(x) \to +\infty$ as $x \to 0$.
By MP, $v_\epsilon(x) \geq 0$ on $B \setminus \{0\}$.
Let ϵ go to zero: $u - v \geq 0$. By replacing u with $(-u)$, we can also conclude $u - v \leq 0$.

Removable Singularity Theorem

Theorem

An isolated singularity of a bounded harmonic function is removable.

Proof: WLOG, u is bounded and harmonic on $\overline{B} \setminus \{0\}$, where $B = B_1(0)$.
Let v be harmonic on B, continuous on $\overline{B}$, with the same boundary values as u.
For $n > 2$, let $v_\epsilon(x) = u(x) - v(x) + \epsilon(|x|^{2-n} - 1)$.
Then $v_\epsilon = 0$ on ∂B, while (u and v are bounded) $v_\epsilon(x) \to +\infty$ as $x \to 0$.
By MP, $v_\epsilon(x) \geq 0$ on $B \setminus \{0\}$.
Let ϵ go to zero: $u - v \geq 0$. By replacing u with $(-u)$, we can also conclude $u - v \leq 0$.
Hence $u = v$.

Positive harmonic functions

Note: We will need our conformal factors to have a sign.

Positive harmonic functions

Note: We will need our conformal factors to have a sign.

Liouville Theorem

Any bounded harmonic function on $\mathbb{R}^n$ is constant.

Positive harmonic functions

Note: We will need our conformal factors to have a sign.

Liouville Theorem

Any bounded harmonic function on $\mathbb{R}^n$ is constant. In fact, any positive harmonic function on $\mathbb{R}^n$ is a constant.

Positive harmonic functions

Note: We will need our conformal factors to have a sign.

Liouville Theorem

Any bounded harmonic function on $\mathbb{R}^n$ is constant. In fact, any positive harmonic function on $\mathbb{R}^n$ is a constant. Thus any harmonic function on $\mathbb{R}^n$ bounded either above or below must be constant.

Positive harmonic functions

Note: We will need our conformal factors to have a sign.

Liouville Theorem

Any bounded harmonic function on $\mathbb{R}^n$ is constant. In fact, any positive harmonic function on $\mathbb{R}^n$ is a constant. Thus any harmonic function on $\mathbb{R}^n$ bounded either above or below must be constant.

Proof: Let $u > 0$ be harmonic on $\mathbb{R}^n$. Use MVP.

Positive harmonic functions

Note: We will need our conformal factors to have a sign.

Liouville Theorem

Any bounded harmonic function on $\mathbb{R}^n$ is constant. In fact, any positive harmonic function on $\mathbb{R}^n$ is a constant. Thus any harmonic function on $\mathbb{R}^n$ bounded either above or below must be constant.

Proof: Let $u > 0$ be harmonic on $\mathbb{R}^n$. Use MVP. Let $R > |x|$.

Positive harmonic functions

Note: We will need our conformal factors to have a sign.

Liouville Theorem

Any bounded harmonic function on $\mathbb{R}^n$ is constant. In fact, any positive harmonic function on $\mathbb{R}^n$ is a constant. Thus any harmonic function on $\mathbb{R}^n$ bounded either above or below must be constant.

Proof: Let $u > 0$ be harmonic on $\mathbb{R}^n$. Use MVP. Let $R > |x|$.

$u(x) - u(0) = \frac{1}{\Omega_n R^n} \left[\int\limits_{B_R(x)} - \int\limits_{B_R(0)} u \, dy \right]$. Since $u > 0$ and $B_R(x) \triangle B_R(0) \subset B_{R+|x|}(0) \setminus B_{R-|x|}(0)$, we have

Positive harmonic functions

Note: We will need our conformal factors to have a sign.

Liouville Theorem

Any bounded harmonic function on $\mathbb{R}^n$ is constant. In fact, any positive harmonic function on $\mathbb{R}^n$ is a constant. Thus any harmonic function on $\mathbb{R}^n$ bounded either above or below must be constant.

Proof: Let $u > 0$ be harmonic on $\mathbb{R}^n$. Use MVP. Let $R > |x|$.

$u(x) - u(0) = \frac{1}{\Omega_n R^n}\left[\int\limits_{B_R(x)} - \int\limits_{B_R(0)} u\, dy\right]$. Since $u > 0$ and $B_R(x) \triangle B_R(0) \subset B_{R+|x|}(0) \setminus B_{R-|x|}(0)$, we have

$|u(x) - u(0)| \leq \frac{1}{\Omega_n R^n} \int\limits_{B_{R+|x|}(0)\setminus B_{R-|x|}(0)} u\, dy =$

$\frac{1}{R^n}((R+|x|)^n - (R-|x|)^n)u(0) = O(R^{-1})$.

Positive harmonic functions

Note: We will need our conformal factors to have a sign.

Liouville Theorem

Any bounded harmonic function on $\mathbb{R}^n$ is constant. In fact, any positive harmonic function on $\mathbb{R}^n$ is a constant. Thus any harmonic function on $\mathbb{R}^n$ bounded either above or below must be constant.

Proof: Let $u > 0$ be harmonic on $\mathbb{R}^n$. Use MVP. Let $R > |x|$.

$u(x) - u(0) = \frac{1}{\Omega_n R^n}\left[\int\limits_{B_R(x)} - \int\limits_{B_R(0)} u\, dy\right]$. Since $u > 0$ and $B_R(x) \triangle B_R(0) \subset B_{R+|x|}(0) \setminus B_{R-|x|}(0)$, we have

$|u(x) - u(0)| \leq \frac{1}{\Omega_n R^n} \int\limits_{B_{R+|x|}(0)\setminus B_{R-|x|}(0)} u\, dy =$

$\frac{1}{R^n}((R+|x|)^n - (R-|x|)^n)u(0) = O(R^{-1})$.

Let $R \to +\infty$.

Positive harmonic functions

Harnack inequality

Ω connected domain, $K \subset \Omega$ compact. There exists $C > 0$ so that for all positive harmonic functions on Ω, for all $x, y \in K$,

$$C^{-1} \leq \frac{u(x)}{u(y)} \leq C.$$

Positive harmonic functions

Harnack inequality

Ω connected domain, $K \subset \Omega$ compact. There exists $C > 0$ so that for all positive harmonic functions on Ω, for all $x, y \in K$,

$$C^{-1} \leq \frac{u(x)}{u(y)} \leq C.$$

Idea: Use MVP. On unit ball B, the following estimate holds for $u \geq 0$:

$$u(x) = \frac{1}{n\Omega_n} \int_{\partial B} \frac{1 - |x|^2}{|x - y|^n} u(y) \, d\sigma_y \leq \frac{1 - |x|^2}{(1 - |x|^n)} u(0).$$

Positive harmonic functions

Harnack inequality

Ω connected domain, $K \subset \Omega$ compact. There exists $C > 0$ so that for all positive harmonic functions on Ω, for all $x, y \in K$,

$$C^{-1} \leq \frac{u(x)}{u(y)} \leq C.$$

Idea: Use MVP. On unit ball B, the following estimate holds for $u \geq 0$:

$$u(x) = \frac{1}{n\Omega_n} \int_{\partial B} \frac{1 - |x|^2}{|x - y|^n} u(y) \, d\sigma_y \leq \frac{1 - |x|^2}{(1 - |x|^n)} u(0).$$

Similarly,

$$u(x) = \frac{1}{n\Omega_n} \int_{\partial B} \frac{1 - |x|^2}{|x - y|^n} u(y) \, d\sigma_y \geq \frac{1 - |x|^2}{(1 + |x|)^n} u(0).$$

Positive harmonic functions

What about positive harmonic functions on $\mathbb{R}^n \setminus \{0\}$?

Positive harmonic functions

What about positive harmonic functions on $\mathbb{R}^n \setminus \{0\}$?

Bôcher's Theorem

If $u > 0$ and harmonic on $B \setminus \{0\}$, then there exists a harmonic function v on B, and a constant $b \geq 0$ so that on $B \setminus \{0\}$,

Positive harmonic functions

What about positive harmonic functions on $\mathbb{R}^n \setminus \{0\}$?

Bôcher's Theorem

If $u > 0$ and harmonic on $B \setminus \{0\}$, then there exists a harmonic function v on B, and a constant $b \geq 0$ so that on $B \setminus \{0\}$,

$$u(x) = \begin{cases} b \log(|x|^{-1}) + v(x) \,, & n = 2 \\ b|x|^{2-n} + v(x) \,, & n > 2 \end{cases}$$

Positive harmonic functions

What about positive harmonic functions on $\mathbb{R}^n \setminus \{0\}$?

Bôcher's Theorem

If $u > 0$ and harmonic on $B \setminus \{0\}$, then there exists a harmonic function v on B, and a constant $b \geq 0$ so that on $B \setminus \{0\}$,

$$u(x) = \begin{cases} b \log(|x|^{-1}) + v(x) \,, & n = 2 \\ b|x|^{2-n} + v(x) \,, & n > 2 \end{cases}$$

Corollary

$n > 2$, $u > 0$ and harmonic on $\mathbb{R}^n \setminus \{0\}$.

Positive harmonic functions

What about positive harmonic functions on $\mathbb{R}^n \setminus \{0\}$?

Bôcher's Theorem

If $u > 0$ and harmonic on $B \setminus \{0\}$, then there exists a harmonic function v on B, and a constant $b \geq 0$ so that on $B \setminus \{0\}$,

$$u(x) = \begin{cases} b \log(|x|^{-1}) + v(x) \,, & n = 2 \\ b|x|^{2-n} + v(x) \,, & n > 2 \end{cases}$$

Corollary

$n > 2$, $u > 0$ and harmonic on $\mathbb{R}^n \setminus \{0\}$. Then there exist numbers $b, c \geq 0$ so that $u(x) = b|x|^{2-n} + c$.

Positive harmonic functions

What about positive harmonic functions on $\mathbb{R}^n \setminus \{0\}$?

Bôcher's Theorem

If $u > 0$ and harmonic on $B \setminus \{0\}$, then there exists a harmonic function v on B, and a constant $b \geq 0$ so that on $B \setminus \{0\}$,

$$u(x) = \begin{cases} b \log(|x|^{-1}) + v(x) \,, & n = 2 \\ b|x|^{2-n} + v(x) \,, & n > 2 \end{cases}$$

Corollary

$n > 2$, $u > 0$ and harmonic on $\mathbb{R}^n \setminus \{0\}$. Then there exist numbers $b, c \geq 0$ so that $u(x) = b|x|^{2-n} + c$.

Proof: On $B \setminus \{0\}$, $u(x) = b|x|^{2-n} + v(x)$. Thus $u(x) - b|x|^{2-n}$ is harmonic on punctured Euclidean space and bounded near the origin. Thus it extends $v(x)$ to a harmonic function on all of $\mathbb{R}^n$.

Positive harmonic functions

What about positive harmonic functions on $\mathbb{R}^n \setminus \{0\}$?

Bôcher's Theorem

If $u > 0$ and harmonic on $B \setminus \{0\}$, then there exists a harmonic function v on B, and a constant $b \geq 0$ so that on $B \setminus \{0\}$,

$$u(x) = \begin{cases} b \log(|x|^{-1}) + v(x) \,, & n = 2 \\ b|x|^{2-n} + v(x) \,, & n > 2 \end{cases}$$

Corollary

$n > 2$, $u > 0$ and harmonic on $\mathbb{R}^n \setminus \{0\}$. Then there exist numbers $b, c \geq 0$ so that $u(x) = b|x|^{2-n} + c$.

Proof: On $B \setminus \{0\}$, $u(x) = b|x|^{2-n} + v(x)$. Thus $u(x) - b|x|^{2-n}$ is harmonic on punctured Euclidean space and bounded near the origin. Thus it extends $v(x)$ to a harmonic function on all of $\mathbb{R}^n$. But $\liminf_{|x| \to +\infty} v(x) \geq 0$.

Positive harmonic functions

What about positive harmonic functions on $\mathbb{R}^n \setminus \{0\}$?

Bôcher's Theorem

If $u > 0$ and harmonic on $B \setminus \{0\}$, then there exists a harmonic function v on B, and a constant $b \geq 0$ so that on $B \setminus \{0\}$,

$$u(x) = \begin{cases} b \log(|x|^{-1}) + v(x) \,, & n = 2 \\ b|x|^{2-n} + v(x) \,, & n > 2 \end{cases}$$

Corollary

$n > 2$, $u > 0$ and harmonic on $\mathbb{R}^n \setminus \{0\}$. Then there exist numbers $b, c \geq 0$ so that $u(x) = b|x|^{2-n} + c$.

Proof: On $B \setminus \{0\}$, $u(x) = b|x|^{2-n} + v(x)$. Thus $u(x) - b|x|^{2-n}$ is harmonic on punctured Euclidean space and bounded near the origin. Thus it extends $v(x)$ to a harmonic function on all of $\mathbb{R}^n$. But $\liminf_{|x| \to +\infty} v(x) \geq 0$. Thus v is constant.

Schwarzschild metric

$g_S = \left(1 + \frac{m}{2|x|^{n-2}}\right)^{4/(n-2)} g_E$ is the spatial Schwarzschild metric of mass m.

Schwarzschild metric

$g_S = \left(1 + \frac{m}{2|x|^{n-2}}\right)^{4/(n-2)} g_E$ is the spatial Schwarzschild metric of mass m.

Since $\frac{1}{|x|^{n-2}}$ is harmonic on $\mathbb{R}^n \setminus \{0\}$, we have $R(g_S) = 0$. It is asymptotically flat with two ends if $m > 0$.

Schwarzschild metric

$g_S = \left(1 + \frac{m}{2|x|^{n-2}}\right)^{4/(n-2)} g_E$ is the spatial Schwarzschild metric of mass m.
Since $\frac{1}{|x|^{n-2}}$ is harmonic on $\mathbb{R}^n \setminus \{0\}$, we have $R(g_S) = 0$. It is asymptotically flat with two ends if $m > 0$.

A simple computation shows (Exercise!)

$$m = \frac{1}{2(n-1)\omega_{n-1}} \lim_{r \to +\infty} \int_{|x|=r} \sum_{i,j=1}^{n} (g_{ij,i} - g_{ii,j}) \nu_e^j \, d\sigma_e.$$

Note $\nu_e^j = \frac{x^j}{|x|}$.

Schwarzschild metric

$g_S = \left(1 + \frac{m}{2|x|^{n-2}}\right)^{4/(n-2)} g_E$ is the spatial Schwarzschild metric of mass m.
Since $\frac{1}{|x|^{n-2}}$ is harmonic on $\mathbb{R}^n \setminus \{0\}$, we have $R(g_S) = 0$. It is asymptotically flat with two ends if $m > 0$.

A simple computation shows (Exercise!)

$$m = \frac{1}{2(n-1)\omega_{n-1}} \lim_{r \to +\infty} \int_{|x|=r} \sum_{i,j=1}^{n} (g_{ij,i} - g_{ii,j}) \nu_e^j \, d\sigma_e.$$

Note $\nu_e^j = \frac{x^j}{|x|}$.

In case $n = 3$, $m = \frac{1}{16\pi} \lim_{r \to +\infty} \int_{|x|=r} \sum_{i,j=1}^{3} (g_{ij,i} - g_{ii,j}) \nu_e^j \, d\sigma_e$.

Schwarzschild metric

$g_S = \left(1 + \frac{m}{2|x|^{n-2}}\right)^{4/(n-2)} g_E$ is the spatial Schwarzschild metric of mass m.
Since $\frac{1}{|x|^{n-2}}$ is harmonic on $\mathbb{R}^n \setminus \{0\}$, we have $R(g_S) = 0$. It is asymptotically flat with two ends if $m > 0$.

A simple computation shows (Exercise!)

$$m = \frac{1}{2(n-1)\omega_{n-1}} \lim_{r \to +\infty} \int_{|x|=r} \sum_{i,j=1}^{n} (g_{ij,i} - g_{ii,j}) \nu_e^j \, d\sigma_e.$$

Note $\nu_e^j = \frac{x^j}{|x|}$.

In case $n = 3$, $m = \frac{1}{16\pi} \lim_{r \to +\infty} \int_{|x|=r} \sum_{i,j=1}^{3} (g_{ij,i} - g_{ii,j}) \nu_e^j \, d\sigma_e$.

The flux integral limit provides the definition of mass/energy for more general asymptotically flat metrics.

Harmonic functions at ∞

Inversion in the unit sphere:

$$x \mapsto x^* = |x|^{-2}x, \quad 0 \mapsto \infty, \quad \infty \mapsto 0.$$

Harmonic functions at ∞

Inversion in the unit sphere:

$$x \mapsto x^* = |x|^{-2}x, \quad 0 \mapsto \infty, \quad \infty \mapsto 0.$$

In $\mathbb{R}^2 \approx \mathbb{C}$, this is $z \mapsto 1/\bar{z}$.

Harmonic functions at ∞

Inversion in the unit sphere:

$$x \mapsto x^* = |x|^{-2}x, \quad 0 \mapsto \infty, \quad \infty \mapsto 0.$$

In $\mathbb{R}^2 \approx \mathbb{C}$, this is $z \mapsto 1/\bar{z}$.

This is a *conformal* transformation.

Harmonic functions at ∞

Inversion in the unit sphere:

$$x \mapsto x^* = |x|^{-2}x, \quad 0 \mapsto \infty, \quad \infty \mapsto 0.$$

In $\mathbb{R}^2 \approx \mathbb{C}$, this is $z \mapsto 1/\bar{z}$.

This is a *conformal* transformation.

Kelvin Transform

For functions u defined on $\Omega \subset \mathbb{R}^n \setminus \{0\}$, define $K[u]$ on Ω^* by

$$K[u](x) = |x|^{2-n}u(x^*).$$

Harmonic functions at ∞

Inversion in the unit sphere:

$$x \mapsto x^* = |x|^{-2}x, \quad 0 \mapsto \infty, \quad \infty \mapsto 0.$$

In $\mathbb{R}^2 \approx \mathbb{C}$, this is $z \mapsto 1/\bar{z}$.

This is a *conformal* transformation.

Kelvin Transform

For functions u defined on $\Omega \subset \mathbb{R}^n \setminus \{0\}$, define $K[u]$ on Ω^* by

$$K[u](x) = |x|^{2-n}u(x^*).$$

Key exercise: u harmonic if and only if $K[u]$ is harmonic.

Harmonic functions at ∞

$K \subset \mathbb{R}^n$ is compact.

Harmonic functions at ∞

$K \subset \mathbb{R}^n$ is compact.

Definition

We say a function u which is harmonic on $\mathbb{R}^n \setminus K$ ("near infinity") is *harmonic at infinity* provided $K[u]$ has a removable singularity at 0.

Harmonic functions at ∞

$K \subset \mathbb{R}^n$ is compact.

Definition

We say a function u which is harmonic on $\mathbb{R}^n \setminus K$ ("near infinity") is *harmonic at infinity* provided $K[u]$ has a removable singularity at 0.

Lemma

$n > 2$. u is harmonic on $\mathbb{R}^n \setminus K$. u is harmonic at infinity if and only if $\lim_{|x| \to +\infty} u(x) = 0$.

Harmonic functions at ∞

$K \subset \mathbb{R}^n$ is compact.

Definition

We say a function u which is harmonic on $\mathbb{R}^n \setminus K$ ("near infinity") is *harmonic at infinity* provided $K[u]$ has a removable singularity at 0.

Lemma

$n > 2$. u is harmonic on $\mathbb{R}^n \setminus K$. u is harmonic at infinity if and only if $\lim_{|x| \to +\infty} u(x) = 0$.

Proof: Exercise! Uses $u(x) = |x^*|^{n-2} K[u](x^*)$.

Harmonic functions at ∞

The Kelvin transform can be used to solve the exterior Dirichlet problem, by inverting solution of the interior Dirichlet problem on ball.

Harmonic functions at ∞

The Kelvin transform can be used to solve the exterior Dirichlet problem, by inverting solution of the interior Dirichlet problem on ball.

The result is a function which is harmonic at infinity.

Harmonic functions at ∞

The Kelvin transform can be used to solve the exterior Dirichlet problem, by inverting solution of the interior Dirichlet problem on ball.

The result is a function which is harmonic at infinity.

Example: $u(x) = (1 - |x|^{2-n})$ is harmonic and $u = 0$ on ∂B. The unique solution of the exterior Dirichlet problem with zero BC's is the zero function. $u(x)$ here is not harmonic at infinity, but $u - 1$ is.

Harmonic functions at ∞

Theorem

For $n > 2$, u harmonic on $\mathbb{R}^n \setminus K$, the following are equivalent.

- u is bounded near infinity
- u is bounded above or below near infinity
- There is a constant c so that $u - c$ is harmonic at infinity
- u has a finite limit as $|x| \to \infty$

Harmonic functions at ∞

Theorem

For $n > 2$, u harmonic on $\mathbb{R}^n \setminus K$, the following are equivalent.

- u is bounded near infinity
- u is bounded above or below near infinity
- There is a constant c so that $u - c$ is harmonic at infinity
- u has a finite limit as $|x| \to \infty$

Proof: Mostly easy. One real step involves using Bôcher's Theorem: if $u > 0$ near infinity, $K[u] > 0$ near the origin, so by Bôcher, there is a c so that $K[u](x) - c|x|^{2-n} = v(x)$ is harmonic near 0. So $K[v](x) = u(x) - c$ is harmonic near infinity.

Spherical harmonic expansion

If v harmonic at infinity,

$$|x^*|^{2-n} v(x) = K[v](x^*) = a_0 + a_i (x^*)^i + \cdots$$

by Taylor expansion of $K[v]$ about $x^* = 0$.

Spherical harmonic expansion

If v harmonic at infinity,

$$|x^*|^{2-n} v(x) = K[v](x^*) = a_0 + a_i (x^*)^i + \cdots$$

by Taylor expansion of $K[v]$ about $x^* = 0$.

Thus $v(x) = a_0 |x^*|^{n-2} + a_i (x^*)^i |x^*|^{n-2} + \cdots = \frac{a_0}{|x|^{n-2}} + \frac{a_i x^i}{|x|^n} + O(|x|^{-n})$.

Spherical harmonic expansion

If v harmonic at infinity,

$$|x^*|^{2-n} v(x) = K[v](x^*) = a_0 + a_i (x^*)^i + \cdots$$

by Taylor expansion of $K[v]$ about $x^* = 0$.

Thus $v(x) = a_0 |x^*|^{n-2} + a_i (x^*)^i |x^*|^{n-2} + \cdots = \frac{a_0}{|x|^{n-2}} + \frac{a_i x^i}{|x|^n} + O(|x|^{-n})$.

One can compute higher-order terms. If you group the Taylor expansion into homogeneous polynomials, each such is harmonic.

Spherical harmonic expansion

If v harmonic at infinity,

$$|x^*|^{2-n} v(x) = K[v](x^*) = a_0 + a_i (x^*)^i + \cdots$$

by Taylor expansion of $K[v]$ about $x^* = 0$.

Thus $v(x) = a_0 |x^*|^{n-2} + a_i (x^*)^i |x^*|^{n-2} + \cdots = \frac{a_0}{|x|^{n-2}} + \frac{a_i x^i}{|x|^n} + O(|x|^{-n})$.

One can compute higher-order terms. If you group the Taylor expansion into homogeneous polynomials, each such is harmonic.

Exercise: In the $n = 3$ case, compute the next term in the expansion of v. What is the dimension of homogeneous harmonic polynomials of degree two in dimension three?

Spherical harmonic expansion

If v harmonic at infinity,

$$|x^*|^{2-n} v(x) = K[v](x^*) = a_0 + a_i (x^*)^i + \cdots$$

by Taylor expansion of $K[v]$ about $x^* = 0$.

Thus $v(x) = a_0 |x^*|^{n-2} + a_i (x^*)^i |x^*|^{n-2} + \cdots = \frac{a_0}{|x|^{n-2}} + \frac{a_i x^i}{|x|^n} + O(|x|^{-n})$.

One can compute higher-order terms. If you group the Taylor expansion into homogeneous polynomials, each such is harmonic.

Exercise: In the $n = 3$ case, compute the next term in the expansion of v. What is the dimension of homogeneous harmonic polynomials of degree two in dimension three?

Remark: The eigenfunctions of Δ_{g_0} on the unit sphere $(\mathbb{S}^n, g_0)$ are given by restrictions of homogeneous harmonic polynomials on $\mathbb{R}^{n+1}$. The first eigenvalue $\lambda_0 = 0$ (constant functions), and the next eigenvalue is $\lambda_1 = n$, with eigenfunctions x^i, $i = 1, \ldots, n+1$ restricted to the sphere.

And now, for something completely different...

Asymptotically Euclidean Metrics and the Positive Mass Theorem

Justin Corvino

Lafayette College
U.S.A.

June 12, 2012

Asymptotically flat metrics

Definition

A complete Riemannian manifold (M^n, g), with an associated symmetric $(0,2)$ tensor K (or $\pi = K - (\mathrm{tr}_g K)g$) is called *asymptotically flat* (or *asymptotically Euclidean*) if there is a compact set $\mathcal{K} \subset M$ so that $M \setminus \mathcal{K}$ equals a disjoint union of asymptotic ends $\bigcup_{j=1}^{k} E_j$, where each asymptotic end E_j is diffeomorphic to $\mathbb{R}^n \setminus \{|x| \leq 1\}$, with asymptotically flat coordinates x^i in which the following decay estimates hold for multi-indices α and β:

Asymptotically flat metrics

Definition

A complete Riemannian manifold (M^n, g), with an associated symmetric $(0,2)$ tensor K (or $\pi = K - (\mathrm{tr}_g K)g$) is called *asymptotically flat* (or *asymptotically Euclidean*) if there is a compact set $\mathcal{K} \subset M$ so that $M \setminus \mathcal{K}$ equals a disjoint union of asymptotic ends $\bigcup_{j=1}^{k} E_j$, where each asymptotic end E_j is diffeomorphic to $\mathbb{R}^n \setminus \{|x| \leq 1\}$, with asymptotically flat coordinates x^i in which the following decay estimates hold for multi-indices α and β:

$$\begin{aligned} |\partial_x^\alpha (g_{ij} - \delta_{ij})(x)| &= O(|x|^{-|\alpha|-q}) \\ \left|\partial_x^\beta K_{ij}(x)\right| &= O(|x|^{-|\beta|-1-q}) \end{aligned}$$

for $|\alpha| \leq \ell + 1$, $|\beta| \leq \ell$, $q > \frac{n-2}{2}$ (so $q > 1/2$ in case $n = 3$).

Example: Schwarzschild

Schwarzschild $m > 0$

$(\mathbb{R}^n \setminus \{0\}, g_S)$ with

$g_S = \left(1 + \frac{m}{2|x|^{n-2}}\right)^{4/(n-2)} g_E$ is asymptotically flat with two ends if $m > 0$.

$q = n - 2$ in this case.

Schwarzschild $m < 0$

Case $n = 3$: $(\mathbb{R}^3 \setminus \{|x| \leq -m/2\}, g_S)$.

Example: Schwarzschild

Schwarzschild $m > 0$

$(\mathbb{R}^n \setminus \{0\}, g_S)$ with
$g_S = \left(1 + \frac{m}{2|x|^{n-2}}\right)^{4/(n-2)} g_E$ is asymptotically flat with two ends if $m > 0$.
$q = n - 2$ in this case.

Schwarzschild $m < 0$

Case $n = 3$: $(\mathbb{R}^3 \setminus \{|x| \leq -m/2\}, g_S)$.
Wait a second, what about positivity of mass?

Example: Schwarzschild

Schwarzschild $m > 0$

$(\mathbb{R}^n \setminus \{0\}, g_S)$ with

$g_S = \left(1 + \frac{m}{2|x|^{n-2}}\right)^{4/(n-2)} g_E$ is asymptotically flat with two ends if $m > 0$.

$q = n - 2$ in this case.

Schwarzschild $m < 0$

Case $n = 3$: $(\mathbb{R}^3 \setminus \{|x| \leq -m/2\}, g_S)$.

Wait a second, what about positivity of mass?

This metric is not complete.

Example: Schwarzschild

Schwarzschild $m > 0$

$(\mathbb{R}^n \setminus \{0\}, g_S)$ with
$g_S = \left(1 + \frac{m}{2|x|^{n-2}}\right)^{4/(n-2)} g_E$ is asymptotically flat with two ends if $m > 0$.
$q = n - 2$ in this case.

Schwarzschild $m < 0$

Case $n = 3$: $(\mathbb{R}^3 \setminus \{|x| \leq -m/2\}, g_S)$.
Wait a second, what about positivity of mass?
This metric is not complete.

Note: if we allow manifolds with boundary, we need to impose something on the boundary: compare Schwarzschild $m > 0$ and $m < 0$.

Example: Schwarzschild

Schwarzschild $m > 0$

$(\mathbb{R}^n \setminus \{0\}, g_S)$ with
$g_S = \left(1 + \frac{m}{2|x|^{n-2}}\right)^{4/(n-2)} g_E$ is asymptotically flat with two ends if $m > 0$.
$q = n - 2$ in this case.

Schwarzschild $m < 0$

Case $n = 3$: $(\mathbb{R}^3 \setminus \{|x| \leq -m/2\}, g_S)$.
Wait a second, what about positivity of mass?
This metric is not complete.

Note: if we allow manifolds with boundary, we need to impose something on the boundary: compare Schwarzschild $m > 0$ and $m < 0$.

For simplicity, let's stick to the boundary-less case.

Example: harmonically flat metrics

Consider an asymptotically flat metric with an end E with asymptotically flat coordinates $x = (x^i)$, so that for $|x| > r_0$, $g = u^4 g_E$, $u \to 1$ at infinity, and $R(g) = 0$, i.e. $\Delta u = 0$.

Example: harmonically flat metrics

Consider an asymptotically flat metric with an end E with asymptotically flat coordinates $x = (x^i)$, so that for $|x| > r_0$, $g = u^4 g_E$, $u \to 1$ at infinity, and $R(g) = 0$, i.e. $\Delta u = 0$.

As we saw earlier, u admits an expansion:

$$u(x) = 1 + \frac{A}{|x|^{n-2}} + \frac{\beta \cdot x}{|x|^n} + \cdots$$

where A is a constant and β is a vector. Note that $q = n - 2$ here.

Example: harmonically flat metrics

Consider an asymptotically flat metric with an end E with asymptotically flat coordinates $x = (x^i)$, so that for $|x| > r_0$, $g = u^4 g_E$, $u \to 1$ at infinity, and $R(g) = 0$, i.e. $\Delta u = 0$.

As we saw earlier, u admits an expansion:

$$u(x) = 1 + \frac{A}{|x|^{n-2}} + \frac{\beta \cdot x}{|x|^n} + \cdots$$

where A is a constant and β is a vector. Note that $q = n - 2$ here.

Note also that derivatives fall off one order faster: $\frac{\partial |x|^{-1}}{\partial x^i} = -|x|^{-3} x^i$.

Example: harmonically flat metrics

Take $n = 3$ for simplicity.

$$u(x) = 1 + \frac{A}{|x|} + \frac{\beta \cdot x}{|x|^3} + \cdots$$

Then we'll see $m = 2A$ (as in Schwarzschild), and if $A \neq 0$, we define $c = \beta/A$: a straightforward computation shows

$\tilde{u}(y) := u(y + c) = 1 + \frac{A}{|y|} + O(|y|^{-3})$.

Example: harmonically flat metrics

Take $n = 3$ for simplicity.

$$u(x) = 1 + \frac{A}{|x|} + \frac{\beta \cdot x}{|x|^3} + \cdots$$

Then we'll see $m = 2A$ (as in Schwarzschild), and if $A \neq 0$, we define $c = \beta/A$: a straightforward computation shows

$\tilde{u}(y) := u(y + c) = 1 + \frac{A}{|y|} + O(|y|^{-3})$.

It makes sense then to identify c as the *center of mass*.

Expansion of scalar curvature

We have $g_{ij} - \delta_{ij} = O(|x|^{-q})$ (and same for inverse metric g^{ij}), $\partial g_{ij} = O(|x|^{-q-1})$ and $\partial^2 g_{ij} = O(|x|^{-q-2})$.

Expansion of scalar curvature

We have $g_{ij} - \delta_{ij} = O(|x|^{-q})$ (and same for inverse metric g^{ij}), $\partial g_{ij} = O(|x|^{-q-1})$ and $\partial^2 g_{ij} = O(|x|^{-q-2})$. Now $\Gamma^k_{ij} = \frac{1}{2} g^{km}(g_{im,j} + g_{mj,i} - g_{ij,m}) = O(|x|^{-q-1})$.

Expansion of scalar curvature

We have $g_{ij} - \delta_{ij} = O(|x|^{-q})$ (and same for inverse metric g^{ij}), $\partial g_{ij} = O(|x|^{-q-1})$ and $\partial^2 g_{ij} = O(|x|^{-q-2})$. Now $\Gamma^k_{ij} = \frac{1}{2} g^{km}(g_{im,j} + g_{mj,i} - g_{ij,m}) = O(|x|^{-q-1})$.

$$\begin{aligned}
\Gamma^k_{ij,\ell} &= \frac{1}{2} g^{km}_{,\ell}(g_{im,j} + g_{mj,i} - g_{ij,m}) + \frac{1}{2} g^{km}(g_{im,j\ell} + g_{mj,i\ell} - g_{ij,m\ell}) \\
&= O(|x|^{-2q-2}) + \frac{1}{2}\delta^{km}(g_{im,j\ell} + g_{mj,i\ell} - g_{ij,m\ell}) \\
&+ \frac{1}{2}(g^{km} - \delta^{km})(g_{im,j\ell} + g_{mj,i\ell} - g_{ij,m\ell}) \\
&= \frac{1}{2}(g_{ik,j\ell} + g_{kj,i\ell} - g_{ij,k\ell}) + O(|x|^{-2q-2}).
\end{aligned}$$

Expansion of scalar curvature

Recall $R(g) = g^{ij}\left(\Gamma^k_{ij,k} - \Gamma^k_{ik,j} + \Gamma^k_{k\ell}\Gamma^\ell_{ij} - \Gamma^k_{j\ell}\Gamma^\ell_{ik}\right)$. Thus
$R(g) = g^{ij}\left(\Gamma^k_{ij,k} - \Gamma^k_{ik,j}\right) + O(|x|^{-2q-2}) = \delta^{ij}\left(\Gamma^k_{ij,k} - \Gamma^k_{ik,j}\right) + O(|x|^{-2q-2})$
because $g - \delta \sim |x|^{-q}$ and $\partial\Gamma \sim |x|^{-q-2}$.

Expansion of scalar curvature

Recall $R(g) = g^{ij}\left(\Gamma^k_{ij,k} - \Gamma^k_{ik,j} + \Gamma^k_{k\ell}\Gamma^\ell_{ij} - \Gamma^k_{j\ell}\Gamma^\ell_{ik}\right)$. Thus
$R(g) = g^{ij}\left(\Gamma^k_{ij,k} - \Gamma^k_{ik,j}\right) + O(|x|^{-2q-2}) = \delta^{ij}\left(\Gamma^k_{ij,k} - \Gamma^k_{ik,j}\right) + O(|x|^{-2q-2})$
because $g - \delta \sim |x|^{-q}$ and $\partial\Gamma \sim |x|^{-q-2}$.

Proposition

Thus using previous expansions,

$$R(g) = \sum_{i,j}(g_{ij,ij} - g_{ii,jj}) + O(|x|^{-2q-2}).$$

Expansion of scalar curvature

Recall $R(g) = g^{ij}\left(\Gamma^k_{ij,k} - \Gamma^k_{ik,j} + \Gamma^k_{k\ell}\Gamma^\ell_{ij} - \Gamma^k_{j\ell}\Gamma^\ell_{ik}\right)$. Thus
$R(g) = g^{ij}\left(\Gamma^k_{ij,k} - \Gamma^k_{ik,j}\right) + O(|x|^{-2q-2}) = \delta^{ij}\left(\Gamma^k_{ij,k} - \Gamma^k_{ik,j}\right) + O(|x|^{-2q-2})$
because $g - \delta \sim |x|^{-q}$ and $\partial\Gamma \sim |x|^{-q-2}$.

Proposition

Thus using previous expansions,

$$R(g) = \sum_{i,j}(g_{ij,ij} - g_{ii,jj}) + O(|x|^{-2q-2}).$$

Note that the error term is in $L^1(E)$: for $q > (n-2)/2$, $2q + 2 > n$.

The mass (energy) integral

Energy

Recall the mass (energy) integral is given by

$$\frac{1}{2(n-1)\omega_{n-1}} \lim_{r\to+\infty} \int_{|x|=r} \sum_{i,j=1}^{n} (g_{ij,i} - g_{ii,j})\nu_e^j \, d\sigma_e.$$

Note $\nu_e^j = \frac{x^j}{|x|}$.

In case $n = 3$, this is $\frac{1}{16\pi} \lim_{r\to+\infty} \int_{|x|=r} \sum_{i,j=1}^{3} (g_{ij,i} - g_{ii,j})\nu_e^j \, d\sigma_e$.

The mass (energy) integral

Energy

Recall the mass (energy) integral is given by

$$\frac{1}{2(n-1)\omega_{n-1}} \lim_{r\to+\infty} \int_{|x|=r} \sum_{i,j=1}^{n} (g_{ij,i} - g_{ii,j})\nu_e^j \, d\sigma_e.$$

Note $\nu_e^j = \frac{x^j}{|x|}$.

In case $n = 3$, this is $\frac{1}{16\pi} \lim_{r\to+\infty} \int_{|x|=r} \sum_{i,j=1}^{3} (g_{ij,i} - g_{ii,j})\nu_e^j \, d\sigma_e$.

Does the limit exist?

The mass (energy) integral

Well, by the divergence theorem,

$$\int_{|x|=r} - \int_{|x|=r_0} \sum_{i,j=1}^{n} (g_{ij,i} - g_{ii,j}) \nu_e^j \, d\sigma_e = \int_{r_0 \le |x| \le r} \sum_{i,j=1}^{n} (g_{ij,ij} - g_{ii,jj}) \, dx$$

$$= \int_{r_0 \le |x| \le r} \sum_{i,j=1}^{n} (R(g) + O(|x|^{-2q-2}) \, dx.$$

The mass (energy) integral

Well, by the divergence theorem,

$$\int_{|x|=r} - \int_{|x|=r_0} \sum_{i,j=1}^{n} (g_{ij,i} - g_{ii,j}) \nu_e^j \, d\sigma_e = \int_{r_0 \le |x| \le r} \sum_{i,j=1}^{n} (g_{ij,ij} - g_{ii,jj}) \, dx$$

$$= \int_{r_0 \le |x| \le r} \sum_{i,j=1}^{n} (R(g) + O(|x|^{-2q-2}) \, dx.$$

Constraints

So if $R(g) \in L^1$, then the limit exists.

The mass (energy) integral

Well, by the divergence theorem,

$$\int_{|x|=r} - \int_{|x|=r_0} \sum_{i,j=1}^{n} (g_{ij,i} - g_{ii,j})\nu_e^j \, d\sigma_e = \int_{r_0 \le |x| \le r} \sum_{i,j=1}^{n} (g_{ij,ij} - g_{ii,jj}) \, dx$$

$$= \int_{r_0 \le |x| \le r} \sum_{i,j=1}^{n} (R(g) + O(|x|^{-2q-2}) \, dx.$$

Constraints

So if $R(g) \in L^1$, then the limit exists.

From the constraint $R(g) - \|K\|_g^2 + H^2 = 2\kappa\rho$, with $K_{ij} = O(|x|^{-q-1})$, we see $R(g) \in L^1(E)$ if and only if $\rho \in L^1(E)$, e.g. vacuum case $\rho = 0$.

The energy-momentum four-vector

Remarks

- The limit can be non-zero even if $R(g) = 0$ (vacuum), e.g. Schwarzschild.

The energy-momentum four-vector

Remarks

- The limit can be non-zero even if $R(g) = 0$ (vacuum), e.g. Schwarzschild.
- One can show (Bartnik, CPAM, 1986) that the mass/energy is independent of AF coordinates.

The energy-momentum four-vector

Remarks

- The limit can be non-zero even if $R(g) = 0$ (vacuum), e.g. Schwarzschild.
- One can show (Bartnik, CPAM, 1986) that the mass/energy is independent of AF coordinates.

Now recall the second constraint equation $\mathrm{div}_g(\pi) = \kappa J$. Now

$$(\mathrm{div}_g \pi)_i = g^{jk}(\pi_{ij,k} - \Gamma^m_{ik}\pi_{mj} - \Gamma^m_{kj}\pi_{im}) = \mathrm{div}_e(\pi) + O(|x|^{-2q-2})$$

The energy-momentum four-vector

Remarks

- The limit can be non-zero even if $R(g) = 0$ (vacuum), e.g. Schwarzschild.
- One can show (Bartnik, CPAM, 1986) that the mass/energy is independent of AF coordinates.

Now recall the second constraint equation $\mathrm{div}_g(\pi) = \kappa J$. Now

$$(\mathrm{div}_g \pi)_i = g^{jk}(\pi_{ij,k} - \Gamma^m_{ik}\pi_{mj} - \Gamma^m_{kj}\pi_{im}) = \mathrm{div}_e(\pi) + O(|x|^{-2q-2})$$

So for $|J| \in L^1(E)$, e.g. $J = 0$, the following limit exists:

$$P_i = \frac{1}{(n-1)\omega_{n-1}} \lim_{r \to +\infty} \int_{|x|=r} \pi_{ij}\nu_e^j \, d\sigma_e.$$

The energy-momentum four-vector

If we let $E = \frac{1}{2(n-1)\omega_{n-1}} \lim_{r\to+\infty} \int_{|x|=r} \sum_{i,j=1}^{n} (g_{ij,i} - g_{ii,j})\nu_e^j \, d\sigma_e$, then E and P fit together to given the *ADM energy-momentum four-vector* of the asymptotically flat end.

The energy-momentum four-vector

If we let $E = \frac{1}{2(n-1)\omega_{n-1}} \lim_{r\to+\infty} \int_{|x|=r} \sum_{i,j=1}^{n} (g_{ij,i} - g_{ii,j})\nu_e^j \, d\sigma_e$, then E and P fit together to given the *ADM energy-momentum four-vector* of the asymptotically flat end.

What do we expect?

The energy-momentum four-vector

If we let $E = \frac{1}{2(n-1)\omega_{n-1}} \lim_{r\to+\infty} \int_{|x|=r} \sum_{i,j=1}^{n} (g_{ij,i} - g_{ii,j})\nu_e^j \, d\sigma_e$, then E and P fit together to given the *ADM energy-momentum four-vector* of the asymptotically flat end.

What do we expect?

For reasonable matter, say corresponding to T satisfying an energy condition, we expect $E \geq 0$, and in $E \geq |P|$, i.e. the energy-momentum vector is future-pointing causal.

The energy-momentum four-vector

If we let $E = \frac{1}{2(n-1)\omega_{n-1}} \lim_{r\to+\infty} \int_{|x|=r} \sum_{i,j=1}^{n} (g_{ij,i} - g_{ii,j})\nu_e^j \, d\sigma_e$, then E and P fit together to given the *ADM energy-momentum four-vector* of the asymptotically flat end.

What do we expect?

For reasonable matter, say corresponding to T satisfying an energy condition, we expect $E \geq 0$, and in $E \geq |P|$, i.e. the energy-momentum vector is future-pointing causal.

This is the content of the *Positive Mass Theorem*.

Positive Mass Theorem

Recall that the dominant energy condition (DEC) implies on the initial data that $\rho \geq |J|$, i.e.

$$\frac{1}{2}(R(g) - \|K\|_g^2 + H^2) \geq |\mathrm{div}_g \pi|.$$

PMT

(M, g, K) is an asymptotically flat initial data set satisfying DEC. Then $E \geq |P|$.

Positive Mass Theorem

Recall that the dominant energy condition (DEC) implies on the initial data that $\rho \geq |J|$, i.e.

$$\frac{1}{2}(R(g) - \|K\|_g^2 + H^2) \geq |\mathrm{div}_g \pi|.$$

PMT

(M, g, K) is an asymptotically flat initial data set satisfying DEC. Then $E \geq |P|$.

PET

(M, g, K) is an asymptotically flat initial data set satisfying DEC. Then $E \geq 0$.

Positive Mass Theorem

Riemannian case

(M, g) is an asymptotically flat initial data set satisfying $R(g) \geq 0$. Then $E \geq 0$.

Positive Mass Theorem

Riemannian case

(M, g) is an asymptotically flat initial data set satisfying $R(g) \geq 0$. Then $E \geq 0$. $E = 0$ implies (M, g) isometric to $(\mathbb{R}^n, g_E)$.

Positive Mass Theorem

Riemannian case

(M, g) is an asymptotically flat initial data set satisfying $R(g) \geq 0$. Then $E \geq 0$. $E = 0$ implies (M, g) isometric to $(\mathbb{R}^n, g_E)$.

Notes

- Riemannian case: Schoen-Yau, 1979 ($3 \leq n \leq 7$), Witten 1981 (spin manifolds).
- PET: Schoen-Yau, 1981 ($n = 3$), Witten (spin); Eichmair ($4 \leq n \leq 7$).
- PMT: Witten, 1981 (spin); Eichmair, Huang, Lee, Schoen, 2011 ($3 \leq n \leq 7$).

Positive Mass Theorem

Riemannian case

(M, g) is an asymptotically flat initial data set satisfying $R(g) \geq 0$. Then $E \geq 0$. $E = 0$ implies (M, g) isometric to $(\mathbb{R}^n, g_E)$.

Notes

- Riemannian case: Schoen-Yau, 1979 ($3 \leq n \leq 7$), Witten 1981 (spin manifolds).
- PET: Schoen-Yau, 1981 ($n = 3$), Witten (spin); Eichmair ($4 \leq n \leq 7$).
- PMT: Witten, 1981 (spin); Eichmair, Huang, Lee, Schoen, 2011 ($3 \leq n \leq 7$).

The rigidity statement in the PMT is: $E = |P|$ only in case M^n is a space-like hypersurface in Minkowski space-time $\mathbb{M}^{n+1}$ with induced metric and second fundamental form g and K.

Positive Mass Theorem

Riemannian case

(M, g) is an asymptotically flat initial data set satisfying $R(g) \geq 0$. Then $E \geq 0$. $E = 0$ implies (M, g) isometric to $(\mathbb{R}^n, g_E)$.

Notes

- Riemannian case: Schoen-Yau, 1979 ($3 \leq n \leq 7$), Witten 1981 (spin manifolds).
- PET: Schoen-Yau, 1981 ($n = 3$), Witten (spin); Eichmair ($4 \leq n \leq 7$).
- PMT: Witten, 1981 (spin); Eichmair, Huang, Lee, Schoen, 2011 ($3 \leq n \leq 7$).

The rigidity statement in the PMT is: $E = |P|$ only in case M^n is a space-like hypersurface in Minkowski space-time $\mathbb{M}^{n+1}$ with induced metric and second fundamental form g and K. This is established in the spin case. Also, in case $E = 0$: Schoen-Yau, Eichmair.

Positive Mass Theorem

Please see arXiv preprint of Eichmair, Huang, Lee and Schoen (2011), and Eichmair (2012) for references and history.

Positive Mass Theorem

Please see arXiv preprint of Eichmair, Huang, Lee and Schoen (2011), and Eichmair (2012) for references and history.

We will sketch a proof the Riemannian case of the PET, at least in case $n = 3$. In this case, the energy E is often just referred to as the mass m, and we'll use the same abuse of notation/miscarriage of justice?

Positive Mass Theorem

Please see arXiv preprint of Eichmair, Huang, Lee and Schoen (2011), and Eichmair (2012) for references and history.

We will sketch a proof the Riemannian case of the PET, at least in case $n = 3$. In this case, the energy E is often just referred to as the mass m, and we'll use the same abuse of notation/miscarriage of justice?

It's OK—in the Riemannian case, can consider $K = 0$, so that $P = 0$, and there is no ambiguity between E and m in this case.

Positive Mass Theorem

Please see arXiv preprint of Eichmair, Huang, Lee and Schoen (2011), and Eichmair (2012) for references and history.

We will sketch a proof the Riemannian case of the PET, at least in case $n = 3$. In this case, the energy E is often just referred to as the mass m, and we'll use the same abuse of notation/miscarriage of justice?

It's OK—in the Riemannian case, can consider $K = 0$, so that $P = 0$, and there is no ambiguity between E and m in this case.

The proof we will sketch blends ideas from the original Schoen-Yau proof, with observations of Bray and Lohkamp on the asymptotics.

Riemannian case

Outline

- Use conformal deformation to reduce to vacuum case $R(g) = 0$.
- Normalize asymptotics I: harmonically flat near infinity
- Simplify: compactify all but one end
- Normalize asymptotics II: Schwarzschild (mass m) near infinity, keeping $R \geq 0$. (Bray)
- If $m < 0$, make the metric EUCLIDEAN near infinity, keeping $R \geq 0$. (Lohkamp)
- Compactify to make a compact manifold with non-flat metric with $R \geq 0$, which is a connected sum with torus. (Lohkamp)
- Argue this metric can be deformed to PSC. Then argue that this is a contradiction—topological obstruction to PSC.
- Rigidity case requires separate argument

Mass under a conformal change

Illustrate in case $n = 3$. Suppose g AF, $q > 1/2$.
$\tilde{g} = u^4 g_{ij}$, where $u = 1 + \frac{A}{|x|} + \cdots$ near infinity (implies drop off in derivatives too).

Mass under a conformal change

Illustrate in case $n = 3$. Suppose g AF, $q > 1/2$.
$\tilde{g} = u^4 g_{ij}$, where $u = 1 + \frac{A}{|x|} + \cdots$ near infinity (implies drop off in derivatives too).

$$\begin{aligned}
(\tilde{g}_{ij,i} - \tilde{g}_{ii,j}) &= u^4(g_{ij,i} - g_{ii,j}) + 4u^3 u_{,i} g_{ij} - 4u^3 u_{,j} g_{ii} \\
&= (g_{ij,i} - g_{ii,j}) + 4(u_{,i}\delta_{ij} - u_{,j}\delta_{ii}) + O(|x|^{-q-2}) \\
&= (g_{ij,i} - g_{ii,j}) + 4(-A\frac{x^i}{|x|^3}\delta_{ij} + A\frac{x^j}{|x|^3}) + O(|x|^{-q-2})
\end{aligned}$$

Mass under a conformal change

Illustrate in case $n = 3$. Suppose g AF, $q > 1/2$.
$\tilde{g} = u^4 g_{ij}$, where $u = 1 + \frac{A}{|x|} + \cdots$ near infinity (implies drop off in derivatives too).

$$\begin{aligned}(\tilde{g}_{ij,i} - \tilde{g}_{ii,j}) &= u^4(g_{ij,i} - g_{ii,j}) + 4u^3 u_{,i} g_{ij} - 4u^3 u_{,j} g_{ii} \\ &= (g_{ij,i} - g_{ii,j}) + 4(u_{,i}\delta_{ij} - u_{,j}\delta_{ii}) + O(|x|^{-q-2}) \\ &= (g_{ij,i} - g_{ii,j}) + 4(-A\frac{x^i}{|x|^3}\delta_{ij} + A\frac{x^j}{|x|^3}) + O(|x|^{-q-2})\end{aligned}$$

The mass integrands are thus related by

$$\sum_{i,j=1}^{3}(\tilde{g}_{ij,i} - \tilde{g}_{ii,j}) = \sum_{i,j=1}^{3}(g_{ij,i} - g_{ii,j}) + 8A\sum_{j=1}^{3}\frac{x^j}{|x|^3} + O(|x|^{-q-2}).$$

The surface integral of the A-term is

$$\frac{1}{16\pi}\int_{|x|=r} 8A\sum_{j=1}^{3}\frac{x^j}{|x|^3}\frac{x^j}{|x|}\,d\sigma_e = \frac{1}{16\pi}\frac{8A}{r^2}\cdot 4\pi r^2 = 2A.$$

Mass under a conformal change

Illustrate in case $n = 3$. Suppose g AF, $q > 1/2$.
$\tilde{g} = u^4 g_{ij}$, where $u = 1 + \frac{A}{|x|} + \cdots$ near infinity (implies drop off in derivatives too).

$$\begin{aligned}(\tilde{g}_{ij,i} - \tilde{g}_{ii,j}) &= u^4(g_{ij,i} - g_{ii,j}) + 4u^3 u_{,i} g_{ij} - 4u^3 u_{,j} g_{ii} \\ &= (g_{ij,i} - g_{ii,j}) + 4(u_{,i}\delta_{ij} - u_{,j}\delta_{ii}) + O(|x|^{-q-2}) \\ &= (g_{ij,i} - g_{ii,j}) + 4(-A\frac{x^i}{|x|^3}\delta_{ij} + A\frac{x^j}{|x|^3}) + O(|x|^{-q-2})\end{aligned}$$

The mass integrands are thus related by

$$\sum_{i,j=1}^{3}(\tilde{g}_{ij,i} - \tilde{g}_{ii,j}) = \sum_{i,j=1}^{3}(g_{ij,i} - g_{ii,j}) + 8A\sum_{j=1}^{3}\frac{x^j}{|x|^3} + O(|x|^{-q-2}).$$

The surface integral of the A-term is

$\frac{1}{16\pi}\int\limits_{|x|=r} 8A\sum_{j=1}^{3}\frac{x^j}{|x|^3}\frac{x^j}{|x|}\,d\sigma_e = \frac{1}{16\pi}\frac{8A}{r^2}\cdot 4\pi r^2 = 2A$. Thus $m(\tilde{g}) = m(g) + 2A$.

Mass under a conformal change

$m(\tilde{g}) = m(g) + 2A.$

As a corollary, this shows the mass of the Schwarzschild metric is indeed $m(g_S) = m(g_E) + 2\frac{m}{2} = 0 + m = m.$

Mass under a conformal change

$m(\tilde{g}) = m(g) + 2A.$

As a corollary, this shows the mass of the Schwarzschild metric is indeed $m(g_S) = m(g_E) + 2\frac{m}{2} = 0 + m = m.$

Note that if $0 < u < 1$, then $A \leq 0$, so that $m(\tilde{g}) \leq m(g)$.

Mass under a conformal change

$m(\tilde{g}) = m(g) + 2A.$

As a corollary, this shows the mass of the Schwarzschild metric is indeed $m(g_S) = m(g_E) + 2\frac{m}{2} = 0 + m = m.$

Note that if $0 < u < 1$, then $A \leq 0$, so that $m(\tilde{g}) \leq m(g)$.

So if the PET works for $\tilde{g}$ (which we might have arranged to be simpler than g in some way), then the PET holds for g.

Laplacian in weighted spaces on AF manifolds

We will use conformal changes of metric to deform the scalar curvature, and so we will want to understand the behavior of the Laplace operator Δ_g, as for $u > 0$:

$$R(u^{4/(n-2)}g) = -\frac{4(n-1)}{n-2}u^{-\frac{n+2}{n-2}}(\Delta_g u - \frac{n-2}{4(n-1)}R(g)u).$$

In particular we will be deforming to zero scalar curvature, so we want to solve $0 = (\Delta_g - \frac{n-2}{4(n-1)}R(g))u$.

Laplacian in weighted spaces on AF manifolds

We will use conformal changes of metric to deform the scalar curvature, and so we will want to understand the behavior of the Laplace operator Δ_g, as for $u > 0$:

$$R(u^{4/(n-2)}g) = -\frac{4(n-1)}{n-2}u^{-\frac{n+2}{n-2}}(\Delta_g u - \frac{n-2}{4(n-1)}R(g)u).$$

In particular we will be deforming to zero scalar curvature, so we want to solve $0 = (\Delta_g - \frac{n-2}{4(n-1)}R(g))u$.

And we want the solution to be positive, $u > 0$, u tends to 1 at infinity (in the relevant end), and if possible, we'd like for u to have at least a partial expansion $u(x) = 1 + \frac{A}{|x|^{n-2}} + O(|x|^{-(n-1)})$ (along with derivative decay).

$n = 3$: $u(x) = 1 + \frac{A}{|x|} + O(|x|^{-2})$

Weighted spaces

(M, g) asymptotically flat. Take $\sigma \geq 1$ a smooth function which in AF coordinates in any end is $\sigma(x) = |x|$.
Define a weighted L^p norm, $p \geq 1$:

$$\|u\|^p_{L^p_{-\tau}} = \int_M (|u|\sigma^\tau)^p \sigma^{-n} \, dv_g.$$

Weighted Sobolev norm

$$\|u\|_{W^{k,p}_{-\tau}} = \sum_{|\gamma| \leq k} \|D^\gamma u\|_{L^p_{-|\gamma|-\tau}}.$$

This gives a Banach space $W^{k,p}_{-\tau}(M, g)$.

(Reference: Bartnik, CPAM, 1986, e.g.)

Weighted spaces

Weighted Hölder spaces are defined analogously. Take $\alpha \in (0, 1)$.

Weighted spaces

Weighted Hölder spaces are defined analogously. Take $\alpha \in (0,1)$.

Let $\sigma(x,y) = \min(\sigma(x), \sigma(y))$, and let $d(x,y)$ be geodesic distance.

$[f]_{\alpha,-\beta} = \sup_{x \neq y} (\sigma(x,y))^{\alpha+\beta} \frac{|f(x)-f(y)|}{(d(x,y)^{\alpha}}$.

Weighted Hölder norm

$$\|f\|_{C^{k,\alpha}_{-\beta}} = \sum_{|\gamma| \leq k} \sup_{x \in M} \left((\sigma(x))^{\beta+|\gamma|} |D^{\gamma} f(x)| \right) + [D^k f]_{\alpha,-\beta-k}.$$

Weighted spaces

A version of the Sobolev embedding is given in the following estimate:

Sobolev Embedding

If $1 \geq k - \frac{n}{p} \geq \alpha > 0$, then $\|u\|_{C^{0,\alpha}_{-\tau}} \leq C\|u\|_{W^{k,p}_{-\tau}}$. If $k > \frac{n}{p}$, $u \in W^{k,p}_{-\delta}$, then $|u(x)| = o(|x|^{-\delta})$.

Both sets of weighted spaces encode decay rates at infinity.

Behavior of Laplacian on weighted spaces

(M, g) AF, with suitable number of derivatives decaying suitably, $k \geq 2$, some range of $p > 1$ (and depends on decay rate of g).

Fredholm property

$\Delta_g : W^{k,p}_{-\delta} \to W^{k-2,p}_{-\delta-2}$ is Fredholm, provided $(-\delta) \notin \{m \in \mathbb{Z} : m \leq 2 - n,\ m \geq 0\}$.

Behavior of Laplacian on weighted spaces

(M, g) AF, with suitable number of derivatives decaying suitably, $k \geq 2$, some range of $p > 1$ (and depends on decay rate of g).

Fredholm property

$\Delta_g : W^{k,p}_{-\delta} \to W^{k-2,p}_{-\delta-2}$ is Fredholm, provided $(-\delta) \notin \{m \in \mathbb{Z} : m \leq 2 - n,\ m \geq 0\}$. (Any comments on these exceptional weights?)

If $\delta \in (0, n-2)$, then Δ_g is invertible. (cf. No Euclidean harmonic functions decay at infinity weaker than $|x|^{2-n}$.) In this case we also have an estimate:

$$\|w\|_{W^{k,p}_{-\delta}} \leq C \|\Delta_g w\|_{W^{k-2,p}_{-\delta-2}}$$

Laplacian on weighted spaces

Conformal Laplacian

For g AF, the map $u \in X^k \to R(g)u \in Y^{k-2}$ is compact in weighted spaces: domain has k derivatives, say, while range has $(k-2)$ derivatives, say.

Laplacian on weighted spaces

Conformal Laplacian

For g AF, the map $u \in X^k \to R(g)u \in Y^{k-2}$ is compact in weighted spaces: domain has k derivatives, say, while range has $(k-2)$ derivatives, say.

Why?

Laplacian on weighted spaces

Conformal Laplacian

For g AF, the map $u \in X^k \to R(g)u \in Y^{k-2}$ is compact in weighted spaces: domain has k derivatives, say, while range has $(k-2)$ derivatives, say.

Why? Ascoli-Arzelà
So $\Delta_g - c(n)R(g)$ will be Fredholm of the same index as Δ_g on the suitable weighted spaces.

Expansion

Let $\delta \in (0, n-2)$. If (M, g) AF, and suppose an equation with Δ_g can be re-written as $\Delta v = F \in W^{2,q}_{-\delta^*-2}$, (for suitable $\tau > 0$, $\delta^* = \delta + \tau$) so that $\Delta : W^{2,q}_{-\delta^*} \to W^{2,q}_{-\delta^*-2}$ is Fredholm, there is a $w \in W^{2,q}_{-\delta^*}$ so that $\Delta(v - w) = 0$ for $|x| > R_0$, say. (Why?)

Then $(v - w)$ goes to zero and is harmonic at infinity, which means $(v - w) = \frac{A}{|x|^{n-2}} + O(|x|^{-(n-1)})$.

Improve estimate on w to get right error term in expansion.

(Cf. Bartnik—weighted Sobolev spaces; N. Meyers, 1963, weighted Hölder spaces)