


presenter: Tony Guttmann, The University of Melbourne

title: An upper bound on the critical exponent for SAWs

problem:

Based on an idea of the late John Hammersley, the following construction provides the basis for a proof that the critical exponent for two-dimensional self-avoiding walks γ , satisfies $\gamma \leq 3$, as usual assuming the existence of such an exponent. The idea relies on the fact that *bridges* are manifestly super-multiplicative, thus if b_n denotes the number of n -step bridges, equivalent up to translation, then $b_n \leq \text{const} \cdot \mu^n$, where μ is the growth constant for SAW. (In fact it is believed that $b_n \sim \text{const} \cdot \mu^n n^{-\frac{7}{16}}$). Bridges are defined as SAW with a unique left-most origin, with end-point being the (not necessarily unique) rightmost point. Thus they can clearly be concatenated without overlap, hence satisfying $b_n \cdot b_m \leq b_{n+m}$, from which the above result follows (also invoking a trivial lower bound).

We define a new class of SAW called *worms*. These are SAW with the same x co-ordinate for the origin and end point. That is to say, their end-points lie on a line parallel to the y -axis. Experimentally, worms are also super-multiplicative, satisfying $w_n \cdot w_m \leq w_{n+m}$, from which would follow a similar bound. However worms can't be concatenated to produce a new worm (in general), so it is not clear how to establish this inequality. Experimentally, we find $w_n \sim \text{const} \cdot \mu^n n^{-\frac{13}{32}}$.



Construction:

In the figure, we have drawn a stylised SAW, and drawn two parallel lattice lines through the origin and the end-point, labelled O and N respectively. Traversing the walk from the origin, A marks the last time the walk contacts the first lattice line, and B marks the first time it contacts the second. Thus the segment OA is a worm, the segment AB is a bridge, and the segment BN is another worm. If the first segment is of length k , the second of length l , then we can write for the total number of SAW of length n , denoted c_n , the following bound:

$$c_n \leq \text{const.} n^2 \cdot \mu^k \cdot \mu^l \cdot \mu^{n-k-l} = \text{const.} \mu^n \cdot n^2.$$

The factor n^2 arises from the possible choices of A and B anywhere on the walk. The missing link in what would be a nice result is our inability to prove that worms are super-multiplicative.